

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/281430117>

# GC/MS Analysis of Essential Oils of *Cymbopogon schoenanthus* and *Origanum majorana* L. Grown in Eastern Algeria

Article in *Asian Journal of Chemistry* · October 2015

DOI: 10.14233/ajchem.2015.18859

CITATION

1

READS

101

4 authors:



**Hadeef Derradji**

Université Kasdi Merbah Ouargla

5 PUBLICATIONS 7 CITATIONS

[SEE PROFILE](#)



**Mohammad Hassan Saidi**

Sharif University of Technology

87 PUBLICATIONS 954 CITATIONS

[SEE PROFILE](#)



**Mohamed Yousfi**

Université Amar Telidji Laghouat

107 PUBLICATIONS 1,786 CITATIONS

[SEE PROFILE](#)



**Moussaoui Yacine**

Université Kasdi Merbah Ouargla

13 PUBLICATIONS 60 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Ph.D. Project [View project](#)



lipase inhibition: obesity treatment [View project](#)



1 **GC/MS Analysis of Essential Oils of *Cymbopogon schoenanthus* and**  
 2 ***Origanum majorana* L. Grown in Eastern Algeria**

3 D. HADEF<sup>1,\*</sup>, M. SAIDI<sup>1</sup>, M. YOUSFI<sup>2</sup> and Y. MOUSSAOUI<sup>3</sup>

4 <sup>1</sup>Laboratoire V.P.R.S, Université de Ouargla, Bp 511 route de Ghardaia, 30000 Ouargla, Algeria

5 <sup>2</sup>Laboratoire des Sciences Fondamentales, Université Amar Telidji de Laghouat, route de Ghardaia, Bp 37G, Laghouat 03000, Algeria

6 <sup>3</sup>Faculté des Mathématique et Science de la Matière, Université de Ouargla, Bp 511 route de Ghardaia.30000 Ouargla, Algeria

7 \*Corresponding author: Fax: +213 29 712627; Tel: +213 29 712627; E-mail: haderadji@yahoo.fr; yacine.moussaoui@gmail.com

(Received: ;

Accepted: )

AJC-0000

8 The chemical compositions of the essential oils obtained from the aerial parts (leaves) of *Cymbopogon schoenanthus* and *Origanum*  
 9 *majorana* L, grown in Eastern Algeria (35°24'15" N and 8°7'27" E), by hydrodistillation was carried out using gas chromatography-mass  
 10 spectrometry (GC-MS) analysis. The components obtained were identified. The most important compound detected in leaves of *Cymbopogon*  
 11 *proxima* L was 4-isopropyl-1-methyl-2-cyclohexen-1-ol (*cis*) (15 %) and most important compound detected in leaves of *Origanum*  
 12 *majorana* L was *p*-menth-1-en-4-ol, (*R*) (21 %).

13 **Keywords:** Essential oil, *Cymbopogon schoenanthus*, *Origanum majorana* L, GC/MS, Hydrodistillation.

## INTRODUCTION

14 The world is rich with natural and unique medicinal plants.  
 15 Medicinal plants are now getting more attention than ever  
 16 because they have potential of myriad benefits to society or  
 17 indeed to all mankind, especially in the line of medicine and  
 18 pharmacological. The medicinal value of these plants lies in  
 19 bioactive phytochemical constituents that produce definite  
 20 physiological action on the human body<sup>1</sup>. Some of the most  
 21 important bioactive phytochemical constituents are alkaloids,  
 22 essential oils, flavonoids, tannins, terpenoid, saponins, phenolic  
 23 compounds and many more<sup>2</sup>. These natural compounds formed  
 24 the foundations of modern prescription drugs as we know  
 25 today<sup>3</sup>. Phytochemical is a natural bioactive compound found  
 26 in plants, such as vegetables, fruits, medicinal plants, flowers,  
 27 leaves and roots that work with nutrients and fibers to act as  
 28 an defense system against disease or more accurately, to protect  
 29 against disease. Phytochemicals are divided into two groups,  
 30 which are primary and secondary constituents; according to  
 31 their functions in plant metabolism. Primary constituents  
 32 comprise common sugars, amino acids, proteins and  
 33 chlorophyll while secondary constituents consists of alkaloids,  
 34 terpenoids and phenolic compounds and many more such as  
 35 flavonoids tannins, etc.<sup>4</sup>.

36 The genus *Origanum* belonging to the *Lamiaceae* family,  
 37 has 38 species that are widespread in the Euro-Siberian and

Irano-Siberian regions. However, most of the species, about 38  
 75 % are concentrates in the Mediterranean periphery, 39  
 especially in the East Mediterranean regions<sup>5-14</sup>. The term 40  
 oregano comes from two Greek words "oros" and "genos" is- 41  
 a` say "radiance of the mountains". *Cymbopogon schoenanthus* 42  
 L Spreng, is an aromatic herb. Fresh young leaves are 43  
 consumed in salads and are used to prepare traditional meat 44  
 recipes. Due to its pleasant aroma and taste it is used to prepare 45  
 an aromatic "tea" that is much appreciated and largely 46  
 consumed in the north of Africa<sup>15</sup>. Besides its use in culinary, 47  
*C. schoenanthus* is also used in folk medicine. Its medicinal 48  
 properties are known from the antiquity, being already 49  
 described by "Pliny the Eldey" in his book *Naturalis Historia*<sup>16</sup>. 50  
 Le Floc'h<sup>17</sup> reports its use for the treatment of rheumatism 51  
 and fever. This author describes also its use as a diuretic, 52  
 insecticide and a poultice to cure dromedary wounds. In the 53  
 South of Tunisia, this plant is also used for the treatment of 54  
 rheumatism and to diminish fever. The plant is particularly 55  
 appreciated for its medicinal action in North Africa and it is 56  
 also used for the anorexia. In the Djanet area (Algeria), it is 57  
 well known for bringing back the appetite. The infusions are 58  
 taken as a diuretic, it cures intestinal troubles and, in the form 59  
 of decoction, it acts against food poisoning and helps also in 60  
 the digestion. Some of the diseases like rheumatism and fever, 61  
 that this plant is used against, can be attributed to the formation 62  
 of free radicals in the biological system<sup>18</sup>. 63

64 The purpose of this study was evaluate the essential oils  
65 of 2 medicinal plants (*Cymbogopogn schoenanthus* and  
66 *Origanum majorana* L) grown in East Algeria (Tebessa region).

### EXPERIMENTAL

67 Samples of the aerial part leaves of (*Cymbogopogn*  
68 *schoenanthus* and *Origanum majorana* L) were collected from  
69 Tebessa region, Eastern Algeria (35°24'15" N and 8°7'27" E).  
70 Plant material were gathered at the flowering stage in March  
71 2014, cut in to little pieces and weighed before the extraction  
72 of volatile compounds. Voucher specimens are deposited in  
73 VPRS Laboratory, University of Ouargla, under the code  
74 Number CP1 and OM1 Respectively, the Fig. 1a and 1b shown  
75 respectively, the picture of *Origanum majorana* L and  
76 *Cymbogopogn schoenanthus*.




Fig. 1. Picture of plants (a): *Origanum majorana* L; (b) *Cymbogopogn schoenanthus*

77 **General procedure:** Conventional hydrodistillation was  
78 carried out with a Clevenger-type apparatus and using samples  
79 of at least 100 g of dried leaves of each plant. The essential oil  
80 of *Cymbogopogn schoenanthus* was obtained after 3 h of  
81 distillation and the yield was 0.40 %. The essential oil of  
82 *Origanum majorana* L was obtained after 4 h of distillation  
83 the yield was 0.60 %. Oils were collected in the vials,  
84 dehydrated with anhydrous sodium sulfate and kept under  
85 refrigeration until being analyzed.

86 **Detection method:** The essential oils were analyzed by  
87 Trace gas chromatograph and a Trace Q mass spectrometer.  
88 These two latter were managed by the dedicated Excalibur

software (all from Thermo Fisher, Rodano MI, Italy) using a  
fused-silica capillary column with an apolar stationary phase  
HP5MSTM (30 m × 0.25 mm × 0.25 μm film thickness). GC-  
MS spectra were obtained using the following conditions:  
carrier gas Helium; flow rate 1.0 mL min<sup>-1</sup>; split 1:20; injection  
volume 0.1 μL; injection temperature 250 °C; oven temperature  
progress from 60 to 280 °C at 2 °C min<sup>-1</sup>; the ionization mode  
used was electronic impact at 70 eV. The identification of  
analytes was carried out by comparing the peak relative  
retention times and mass spectra.

### RESULTS AND DISCUSSION

Fig. 2, represent the chromatogram GC/MS of essential oil from  
*Cymbogopogn schoenanthus* in scan mode, while the  
Fig. 3 represent the chromatogram GC/MS of essential oil from  
*Origanum majorana* L in scan mode.




Fig. 2. GC/MS chromatogram of essential oil from *Cymbogopogn schoenanthus*




Fig. 3. GC/MS chromatogram of essential oil from *Origanum majorana* L.

103 The Table-1, regroup the most abundant compounds of  
 104 essential oil composition, twenty one compounds were  
 105 identified as majorities compounds. The compounds identified  
 106 were included in four families: monoterpenes hydrocarbons;  
 107 oxygenated monoterpenes; sesquiterpenes hydrocarbons and  
 108 oxygenated sesquiterpenes. Oxygenated monoterpenes was the  
 109 most abundant family, which represent 49 % (Fig. 4) and 4-  
 110 isopropyl-1-methyl-2-cyclohexen-1-ol (*cis*) was the most  
 111 abundant compound when represent 15 % of the total  
 112 compounds identified.

113 For the essential oil composition obtained from *Origanum*  
 114 *majorana* L, the monoterpenes hydrocarbons and oxygenated  
 115 monoterpenes were the majorities family, they represent 90 %  
 116 of the total, while the sesquiterpenes hydrocarbons and

oxygenated sesquiterpenes families represent together 10 % 117  
 (Fig. 5). Among the essential oil composition, *p*-menth-1-en- 118  
 4-ol, (R) was the most abundant compounds with 21 % of the 119  
 total essential oil compounds identified. 120

The composition of essential oil compounds of the areal 121  
 parts from two plants does not include many very volatile 122  
 compounds; this fact will be in relation with oven temperature 123  
 program of gas chromatography. Since that, the temperature 124  
 program started at 60 °C and at this temperature does not 125  
 promote the very volatile compounds in the chromatogram. 126  
 Other analyse will be focused of identification of very volatile 127  
 compounds from two plants in the future using head space/ 128  
 solid phase microextraction (HS/SPME). 129

TABLE-1  
 SEMI-QUANTITATIVE COMPOSITION OF ESSENTIAL OIL FROM *Cymbopogon schoenanthus*

No.	Retention time (min)	Compounds	%
1	8.63	2-Carene	7
2	9.00	$\alpha$ -Phellandrene	3
3	9.44	<i>p</i> -Mentha-1,4(8)-diene	1
4	9.89	<i>p</i> -Cymene	2
5	10.02	d-Limonene	5
6	10.16	$\beta$ -Phellandrene	5
7	15.40	4-Isopropyl-1-methyl-2-cyclohexen-1-ol ( <i>cis</i> )	15
8	16.49	4-Isopropyl-1-methyl-2-cyclohexen-1-ol ( <i>trans</i> )	13
9	16.76	Camphor{1R,4R}	1
10	19.65	<i>p</i> -Menth-1-en-3-ol, <i>cis</i>	8
11	20.51	<i>p</i> -Menth-1-en-3-ol, <i>trans</i>	9
12	22.23	<i>p</i> -Menth-4(8)-en-3-one	2
13	23.30	<i>p</i> -Menth-1-en-3-one	1
14	25.69	Carvacrol	1
15	31.11	Cyclohexene, 2,4-diisopropenyl-1-methyl-1-vinyl-(1S, 2R, 4R)	3
16	32.86	6-Chamigrene	1
17	37.22	Isolongifolan-8-ol	2
18	38.73	$\gamma$ -Cadinene	2
19	41.09	Cyclohexanemethanol, 4-ethylene- $\alpha,\alpha$ , 4-trimethyl-3-(1-methylethenyl)-, [1R(1 $\alpha$ ,3 $\alpha$ , 4 $\beta$ )]-	10
20	47.18	2-Naphtalenemethanol, decahydro- $\alpha,\alpha$ , 4a-trimethyl-8-methylene-, [2R(2 $\alpha$ ,4 $\alpha$ , 8 $\alpha\beta$ )]-	9
21	57.47	1,7,7-Trimethylbicyclo(2,2,1)-hept-2-yl 3-methylenecyclopentacarboxylate	< 1

TABLE-2  
 SEMI-QUANTITATIVE COMPOSITION OF ESSENTIAL OIL FROM *Origanum majorana* L.

No.	Retention time (min)	Compounds	%
1	7.59	$\beta$ -Phellandrene	6
2	9.45	$\alpha$ -Terpinene	15
3	11.67	$\tau$ -Terpinene	19
4	12.16	4-Isopropogenyl-1-methylcyclohexanol	3
5	12.80	<i>p</i> -Mentha-1,4(8)-diene	5
6	13.90	Terpineol ( <i>cis</i> )	11
7	15.16	4-Isopropyl-1 methyl-2-cyclohexen-1-ol	1
8	16.76	Camphor{1R,4R}	3
9	18.64	<i>p</i> -Menth-1-en-4-ol, (R)	21
10	19.58	<i>p</i> -Menth-1-en-8-ol, (S)	8
11	20.32	<i>p</i> -Menth-1-en-3-ol <i>trans</i>	1
12	22.49	Acetic acid linalool ester	1
13	24.78	Bornyl acetate	< 1
14	25.69	<i>p</i> -Menth-1-en-4-ol acetate	1
15	32.88	Caryophyllene	1
16	37.57	<i>o</i> -Menth-8-ene, 4-isopropylidene-1-vinyl	< 1
17	38.39	1,5-Heptadiene, 6-methyl-2-(4-methyl-3-cyclohexen-1-yl)-, (S)	< 1
18	42.56	Spathulenol	< 1
19	44.01	Carotol	< 1
20	47.79	Ar-Tumenone	< 1




Fig. 4. Composition of essential oil by family from *Cymbopogon schoenanthus*




Fig. 5. Composition of essential oil by family from *Origanum majorana* L.

130 The reported data literature for the content of the main  
 131 components of *Origanum majorana* essential oils shows  
 132 different main components according to the origin of the plant.  
 133 Essential oil of *Origanum majorana* was rich in cymyl  
 134 compounds (carvacrol 77 %) of *Origanum majorana* from  
 135 Argentina<sup>19</sup>, 1,8-cineole (33.5-50.9 %) from Italy<sup>20,21</sup>, Linalool  
 136 (15.5-37.8 %) from Finland<sup>22</sup>, terpinen-4-ol (20.0-55.1 %)   
 137 from Austria<sup>23</sup> and Egypt<sup>24</sup> and  $\gamma$ -terpinene (0.5-14.1 %)   
 138 from Germany<sup>25</sup> and Hungary<sup>26</sup> and *p*-cymene (0.0-11.3 %)   
 139 from Poland<sup>27</sup>, Reunion island<sup>28</sup> and Tunisia<sup>29</sup>. In our study, the  
 140 essential oil of *Origanum majorana* L was rich in *p*-menth-1-  
 141 en-4-ol, (R).

142 The previous study of essential oil of *Cymbopogon*  
 143 *schoenanthus* from Togo, indicate that the oil was rich in  
 144 piperitone (68 %) and 2-carene (21 %) <sup>30</sup>, while the essential  
 145 oil from *Cymbopogon schoenanthus* in this study was rich in  
 146 4-Isopropyl-1-methyl-2-cyclohexen-1-ol (*cis*) (15 %).

## 147 Conclusion

148 The study of the essential oil of the aerial parts of  
 149 *Cymbopogon schoenanthus* and *Origanum majorana* L.  
 150 allowed to identify the most abundant compounds of essential  
 151 oil composition.

Twenty one and twenty compounds were identified as 152  
 majorities compounds of essential oil respectively from 153  
*Cymbopogon schoenanthus* and *Origanum majorana* L. 154

4-Isopropyl-1-methyl-2-cyclohexen-1-ol (*cis*) was the 155  
 most abundant compound when represent 15 % of the total 156  
 compounds identified of oil essential from *Cymbopogon* 157  
*schoenanthus*. While *p*-menth-1-en-4-ol, (R) was the most 158  
 abundant compounds with 21 % of the total essential oil comp- 159  
 ounds identified of oil essential from *Origanum majorana* L. 160

## REFERENCES

1. A.C. Akinmoladun, E.O. Ibukun, E. Afor, E.M. Obuotor and E.O. Farombi, *Sci. Res. Essays*, **2**, 163 (2007).
2. H.O. Edeoga, D.E. Okwu and B.O. Mbaebie, *Afr. J. Biotechnol.*, **4**, 685 (2005).
3. S.H. Goh, C.H. Chuah, J.S.L. Mok and E. Soepadmo, Pelanduk Publication. Kaula Lumpur, Malaysia (1995).
4. D. Krishnaiah, R. Sarbatly and A. Bono, *Biotechnol. Mol. Biol. Rev.*, **1**, 97 (2007).
5. J.H. Ietswaart, Leiden University Press, la Haye (1980).
6. A.M. Janssen, N.L.J. Chin, J.J.C. Scheffer and A.B. Svendsen, *Pharm. Weekbl. Sci.*, **8**, 289 (1986).
7. R.J.W. Lambert, P.N. Skandamis, P. Coote and G.J.E. Nychas, *J. Appl. Microbiol.*, **91**, 453 (2001).
8. A. Morris, A. Khettry and E.W. Seitz, *J. Am. Oil Chem. Soc.*, **56**, 595 (1979).
9. M. Nevas, A.R. Korhonen, M. Lindstrom, P. Turkki and H. Korkela, *J. Food Prot.*, **67**, 199 (2004).
10. B. Ouattara, R.E. Simard, R.A. Holley, G.J.-P. Piette and A. Bégin, *Int. J. Food Microbiol.*, **37**, 155 (1997).
11. M. Oussalah, S. Caillet, L. Saucier and M. Lacroix, *Food Contr.*, **18**, 414 (2007).
12. R. Piccaglia, M. Marotti, E. Giovanelli, S.G. Deans and E. Eaglesham, *Ind. Crops Prod.*, **2**, 47 (1993).
13. A. Remmal, T. Bouchikhi and M. Rhayour-Ettaybi, *J. Essent. Oil Res.*, **5**, 179 (1993).
14. G. Ruberto, M.T. Barratta, M. Sari and M. Kaabeche, *Flavour Fragrance J.*, **17**, 251 (2002).
15. IUCN A guide to medicinal plants in North Africa. ISBN: 2, 8317-0893-1, p. 256 (2005).
16. P. L'Ancien, ed., d'Emile Littré, Paris (1848-1850).
17. E. Le Floc'h, Ministère de l'Enseignement Supérieure et de la Recherche Scientifique, Tunis (1983).
18. K. Bauerova and A. Bezek, *Gen. Physiol. Biophys.*, **18**, 15 (1999).
19. D.M. Maestri, J.A. Zygodlo, A.L. Lamarque, D.O. Labuckas and C.A. Guzmán, *Grasas Aceites*, **47**, 397 (1996).
20. I. Camele, V. De Feo, L. Altieri, E. Mancini, L. De Martino and G. Luigi Rana, *J. Med. Food*, **13**, 1515 (2010).
21. L.F.R. de Almeida, F. Frei, E. Mancini, L. De Martino and V. De Feo, *Molecules*, **15**, 4309 (2010).
22. S. Hälvä, *J. Agric. Sci. Finland*, **59**, 31 (1987).
23. E.H. Koschier, K.A. Sedy and J. Novak, *Crop Prot.*, **21**, 419 (2002).
24. A.E. Edris, A. Shalaby and H.M. Fadel, *Flavour Fragrance J.*, **18**, 345 (2003).
25. J. Richter and I. Schellenberg, *Anal. Bioanal. Chem.*, **387**, 2207 (2007).
26. E. Vági, B. Simándi, A. Suhajda and E. Héthelyi, *Food Res. Int.*, **38**, 51 (2005).
27. A. Dawidowicz and E. Rado, *J. Pharm. Biomed. Anal.*, **52**, 79 (2010).
28. R.R. Vera and J. Chané-Ming, *Food Chem.*, **66**, 143 (1999).
29. I.H. Sellami, E. Maamouri, T. Chahed, W.A. Wannes, M.E. Kchouk and B. Marzouk, *Ind. Crops Prod.*, **30**, 395 (2009).
30. K. Koba, K. Sanda, C. Raynaud, Y.A. Nenonene, J. Millet and J.P. Chaumont, *Ann. Med. Vet.*, **148**, 202 (2004).