

The status of Algeria's sea cucumbers and their illegal trade

Karim Mezali^{1*} and Farah Slimane-Tamacha¹

Abstract

In Algeria, sea cucumber fishing is mainly done by divers and gleaners who target all holothurian species along the coast, especially the ones found in shallow water. To better understand how this fishery is organised, a survey took place from August to September 2018. Face-to-face interviews with fishers (gleaners and divers), members of coastal communities and local traders were carried out to gather detailed information on the fishery. In addition, an online survey was developed to allow stakeholders to respond anonymously. Seventy-eight responses were received from fishers and middlemen who were very familiar with the fishing and trade. These responses were received from 14 different *wilayas* (administrative areas) spread along the Algerian coast. The *wilaya* of Tipaza was found to be the most active in the fishery, followed by Mostaganem, Jijel and Algiers. Sea cucumber processing methods, sales and prices are also discussed, as well as, given the rapid development of the sea cucumber fishery, the urgent need for its effective management.

Key words: Sea cucumber, survey, illegal fishing, Algerian coastal waters

Introduction

Sea cucumbers are in high demand in Asian markets, mainly as a dried product called beche-de-mer (Purcell et al. 2018). Overfishing of sea cucumbers in many tropical countries has led to the capture of new target species from the Mediterranean Sea and the northeast Atlantic Ocean, such as *Holothuria tubulosa*, *H. poli*, *H. sanctori*, *H. arguinensis* and *Parastichopus regalis* (Gonzalez-Wangüemert et al. 2016; Neghli and Mezali 2019). The sea cucumber fishery is a minor activity in Algeria. Basic information on this fishery – such as the number of stakeholders involved, volumes caught and traded, and natural stock estimates – is lacking.

In Algeria, the law allows the collection of sea cucumbers, but not for commercial purposes (Order of 16 July 2008). The export of sea cucumbers is, therefore, illegal, which explains why so little information on the fishery is available. Nevertheless, illegal exports began in 2013 (Neghli and Mezali 2019) and this occurred following a growing Chinese presence in the country, which started in the early 2000s. This non-native population, fond of seafood and with contacts in China, and the main market for processed sea cucumbers, generated a high local demand for sea cucumbers. In response, a fishery developed along the entire Algerian coast before any monitoring or management plan could be put in place. The objective of the study described here was to collect information on the state of exploitation (catches, sales and processing) of sea cucumbers over 14 Algerian coastal *wilayas*.

Methods – Study area and survey

Our study was spread over 14 coastal *wilayas* (administrative divisions) along the Algerian coast: Tlemcen, Ain Timouchent, Oran and Mostaganem in the west; Chlef, Tipaza, Algiers, Boumerdes and Tizi-Ouzou in the middle; and Bejaia, Jijel, Skikda, Annaba and El Taref in the east. Our survey took place between July and September 2018 through face-to-face interviews with people involved in the sea cucumber trade (local traders, fishers and coastal communities). During our survey, we faced two main difficulties: 1) the reluctance or categorical refusal of the majority of fishers to answer our questions, and 2) the non-cooperation of the middlemen involved in the processing and export of the final product (i.e. beche-de-mer). For these reasons, we also developed an online survey in Arabic, the most spoken language in Algeria, which allowed people to respond anonymously.² It included questions such as: “Do you have a preference for a specific species of sea cucumber?” Responses allowed us to determine the most targeted species of sea cucumbers.

Results

Characteristics of the sea cucumber fishery in Algeria

In shallow infralittoral areas, holothurians are most often collected by hand and by free diving on soft or hard substratum. When sea cucumbers are no longer available at these shallow

¹ Protection, Valorisation of Marine and Coastal Resources and Molecular Systematics Laboratory, Department of Marine Sciences and aquaculture, Faculty of Natural Sciences and Life, Abdelhamid Ibn Badis University-Mostaganem, 27000, PO Box 227, Algeria.

² <https://forms.gle/71VWkx9cQPBBQ9XT9>

* Corresponding author: karim.mezali@univ-mosta.dz

depths, fishers use small motorised boats to collect them at greater depths, using direct techniques (scuba diving), and/or indirect methods such as collecting bycatch (starfish) from fishers using demersal gill nets. In Algeria, professional and amateur fishers have long used sea cucumbers as fishing bait (Francour 1997; Mezali 1998). During our survey, we noted that longline fishers tended to replace their fishing bait of small pelagic fishes (e.g. sardines, horse mackerels, *Sardinella* spp.) and cephalopods (mainly squid) with juvenile, or parts of adult, sea cucumbers, due to the rising prices of these small pelagic fishes and cephalopods. We also noted that anglers prefer to use sea cucumbers to catch Sparidae (mainly the sea bream *Sparus aurata*), which are much appreciated in Algerian restaurants.

Most targeted sea cucumber species and sizes

All local sea cucumber species are targeted. Out of 78 respondents, 30% declared a preference for *Holothuria poli*, 27% for *H. tubulosa*, 25% for *H. arguinensis* and 18% for *H. sanctori* (Fig. 1).

Holothuria poli and *H. tubulosa* (Fig. 3A) are the most abundant sea cucumbers in Algerian waters. *H. arguinensis* (Fig. 3B) is a species originating from the Atlantic Ocean but recently reported as occurring in Algerian coastal waters by Mezali and Thandar (2014). *H. arguinensis* is highly valued for its thick body wall. *H. sanctori* (Fig. 3C and D) releases, soon after capture, Cuvierian tubules, which are sticky white threads that function as a defence mechanism. Once ejected, these tubules instantaneously stick to anything (such as hands of a collector) (Flammang 2006). This may explain why it is the least preferred of the four main species.

All sizes of sea cucumbers are targeted: small and medium-sized individuals are used as fishing bait, while medium and large individuals are sold and processed into beche-de-mer.




Figure 1. Fishers' preference for targeted species (in percentage).

Catch rates in the different wilayas

Sea cucumbers are fished all along the Algerian coast, but according to our survey, the best catch rates are obtained along the central coast (Tipaza and Algiers), followed by Jijel in the east and Chlef in the south (Fig. 2).




Figure 2. Catch rates (number of sea cucumbers collected per fishing trip) in each of the 14 surveyed wilayas.

Processing of sea cucumber catches in Algeria

Harvested sea cucumbers are usually kept in basins and stored in large quantities before their transformation into beche-de-mer by processors (Fig. 3).




Figure 3. The processing of sea cucumbers: A) Individuals of different species kept alive in a basin. The processor is holding a specimen of *Holothuria tubulosa*; B) *H. sanctori*; C) Adult *H. arguinensis*; D) *H. arguinensis*. (Photos taken by a sea cucumber processor in Algiers, 2019)

According to our survey, 75% of harvested sea cucumbers are sold fresh to processors, but also to anglers as bait, and to Chinese nationals who reside in Algeria. Fishers are ignorant of the fate of these captured and sold animals. The remaining 25% is directly processed by fishers into beche-de-mer, using processing methods tailored to the targeted export market requirements. The most commonly used sea cucumber processing method involves: 1) evisceration, which can be done by the fishers themselves or by processors; 2) cooking in seawater for two hours; and 3) sun drying for six consecutive days or more. The duration of this last step depends on the season and geographical location. Some processors also salt and/or smoke sea cucumbers as a method of conserving them before drying and selling. These traditional methods of processing sea cucumber into beche-de-mer are the ones used in Algeria.

Sea cucumber trade and sale in Algeria

According to our survey, the selling price of sea cucumbers varies from one *wilaya* to another. Fresh animal prices vary from DZD³ 200–700/kilo (EUR 1.5–5.9/kilo), while dried specimens (beche-de-mer) range in price from DZD 4800–12,000/kilo (EUR 36–90/kilo). Price also depends on demand for the product, its size, its availability and its destination.

Regulation of the sea cucumber fishery in Algeria

Algeria should be inspired by countries in the Pacific Islands region, where minimum sizes and seasonal closures of the fishery are commonly used measures to regulate the sea cucumber fishery (Kinch et al. 2008). In the hope of regulating sea cucumber exploitation, preventing overfishing and avoiding problems with customs (at the trade port), the administration in charge of the fisheries sector published two orders in the official newspaper of the Algerian Republic. The first (Order of 16 July 2008) to allow the collection of sea cucumbers by gleaning and by diving, but not to exploit them commercially because they are not listed in the National Plan of the Marine Biological Resources Regulation, which defines the minimum sizes allowed for commercially exploited resources (Neghli and Mezali 2019); and the second (Decree of 29 Ramadhan 1440, corresponding to 3 June 2019) to set a yearly 46-day closing period (from 1 August to 15 September) for sea cucumber fishing.

Unfortunately, the administration in charge of the fisheries sector does not collaborate with scientists and ignores the fisheries data they have already collected, as well as the scientific work (mainly on reproductive biology) they have already carried out and published in international journals (Neghli et al. 2013; Mezali and Soualili 2014; Mezali et al. 2014; Slimane-Tamacha et al. 2019). The 46-day closure, for example, only partially covers the breeding season of the various sea cucumber species present in Algerian waters.

Discussion

In Algeria, sea cucumber fishing is carried out mainly by hand in infralittoral waters, especially in shallow habitats of seagrass (*Posidonia oceanica*) meadows (Mezali 1998, 2004, 2008). This ease of access allows for intense fishing pressure in these areas, often leading to a local collapse of the most common sea cucumber populations. Fishers who collect sea cucumbers mostly sell them to processors or exporters. However, a survey conducted in the Algerian central region (Neghli and Mezali 2019) revealed the existence of a nascent network of fishers who process sea cucumbers into beche-de-mer by themselves. The depletion of sea cucumbers in shallow water areas of some Algerian *wilayas* has led to exploration of deeper zones. This spatial expansion has already been described for many tropical sea cucumber fisheries (e.g. Maldives, Philippines and Sri Lanka), where sea cucumbers catches first started manually near the coast before being done by diving or even trawling as shallow stocks diminished (Anderson et al. 2011).

According to the interviews we conducted with fishers in different Algerian *wilayas*, the sales price of fresh and dried sea cucumbers varies greatly from one *wilaya* to another and from one fisher to another. In some cases, it was difficult for us to know the exact selling price and destination of the final product due to the withholding of information by some traders. In general, it is estimated that, despite being the main actors in this illegal fishery, fishers are the ones within the market chain who benefit the least from the trade. The dry weight of the most properly processed sea cucumbers is 5–12% of their wet weight (Choo 2008). According to the information provided in our survey, the price of a dry processed sea cucumber is generally 17–24 times higher than the price of fresh (wet) sea cucumber. The majority of fishers do not master all of the steps of processing sea cucumbers into beche-de-mer. In fact, the only part of processing they master is evisceration, which brings very little added value to the product.

Furthermore, the lack of control over actual catches makes it difficult to develop analytical models to support the management of sea cucumber stocks. It is important to adopt a well-defined management plan to preserve this marine resource.

In order to manage sea cucumber resources in Algeria, several important measures must be taken. The development of guidelines for the management of natural stocks of sea cucumbers, and the monitoring of natural stock replenishment, are a priority. The management plan should also include the following measures:

- 1) Prohibit the collection of sea cucumbers during the breeding season of each species; the closing period should be extended (to June–September) or shifted (to July–August, rather than August–September) (Neghli et al. 2013; Mezali and Soualili 2014; Mezali et al. 2014; Slimane-Tamacha et al. 2019).

³ DZD = Algerian dinars (DZD 100 = EUR 0.76 = USD 0.83, as of 12 February 2020)

- 2) Introduce a quota system.
- 3) Define minimum sizes for each species.
- 4) Create permanent protected areas in different *wilayas* along the Algerian coastline.
- 5) Issue fishing licenses to sea cucumber fishers.
- 6) Draft detailed datasheets, and make it mandatory for fishers, processors and traders to fill in and return these sheets to local fisheries authorities.
- 7) Train beche-de-mer processors to upgrade their skills.
- 8) Train all stakeholders who participate in the protection of coastal marine resources (e.g. coastguard officials, custom officers, fisheries directorate agents) in the proper identification of sea cucumber species exploited in Algeria, both in their live and dried forms, using illustrated guides and manuals.
- 9) Monitor all stages of the commercial sea cucumber trade in the country.

Acknowledgements

The authors are extremely grateful to all the fisheries colleagues who contributed to the sharing of the questionnaires and facilitated contact with sea cucumber fishers. We also thank the divers and other professional fishers for their cooperation and trust. The first author thanks the editors for their constructive comments.

References

- Anderson S.C., Flemming J.M., Watson R., Lotze H.K. and Bograd S.J. 2011. Rapid global expansion of invertebrate fisheries: Trends, drivers, and ecosystem effects. *Plos One* 6, e14735.
- Choo P.S. 2008. Population status, fisheries and trade of sea cucumbers in Asia. p. 81–118. In: Toral-Granda V., Lovatelli A. and Vasconcellos M. (eds). *Sea cucumbers: A global review of fisheries and trade*. Fisheries and Aquaculture. Technical Paper 516. Rome: Food and Agriculture Organization of the United Nations. 317 p.
- Flammang P. 2006. Adhesive secretions in echinoderms: An overview. P. 183–206. In: Smith A.M. and Callow J.A. (eds). *Biological adhesives*. Geidelberg, Germany: Springer-Verlag.
- Francour P. 1997. Predation of holothurians: A literature review. *Invertebrate Biology* 116(1):52–60.
- Gonzalez-Wangüemert M., Valente S., Henriques F., Domínguez-Godino J. and Serrao E. 2016. Setting preliminary biometric baselines for new target sea cucumbers species of the NE Atlantic and Mediterranean fisheries. *Fisheries Research* 179: 57–66.
- Kinch J., Purcell S., Uthicke S. and Friedman K. 2008. Population status, fisheries and trade of sea cucumbers in the Western Pacific. p. 7–55. In: Toral-Granda V., Lovatelli A. and Vasconcellos M. (eds). *Sea cucumbers: A global*

review on fisheries and trade. Fisheries and Aquaculture Technical Paper. No. 516. Rome: Food and Agriculture Organization of the United Nations.

- Mezali K. 1998. Contribution à la systématique, la biologie, l'écologie et la dynamique de cinq espèces d'holothuries aspidochirotes (*Holothuria* (*H.*) *tubulosa*, *H.* (*L.*) *poli*, *H.* (*H.*) *stellati*, *H.* (*P.*) *forskali* and *H.* (*P.*) *sanctori*) inféodées aux herbiers à *Posidonia oceanica* (L.) de la péninsule de Sidi-Fredj. Mémoire de Magister. Ecole National des Sciences de la Mer et de L'aménagement du Littoral. Alger, Algérie. 192 p.
- Mezali K. 2004. Micro-répartition des holothuries aspidochirotes au sein de l'herbier de Posidonies de la presqu'île de Sidi-Fredj – Algérie. Rapports P.V. Commission International pour l'Exploration Scientifique de la Mer Méditerranée vol. 37. 534 p.
- Mezali K. 2008. Phylogénie, systématique, dynamique des populations et nutrition de quelques espèces d'holothuries aspidochirotes (Holothuroidea: Echinodermata) inféodées aux herbiers de Posidonies de la côte algéroise. Thèse de Doctorat d'état. Institut des Sciences Biologiques/Université des Sciences et de la Technologie Houari Boumediene, Alger, Algérie. 208 p.
- Mezali K. and Soualili D.L. 2014. Reproductive cycle of the sea cucumber *Holothuria tubulosa* (Holothuroidea: Echinodermata) in the southwestern Mediterranean Sea. Communication orale. Congrès International sur les "Les Estuaires et Aires Marines Côtières Protégées - ECPA 2014" Izmir-Turquie, 4–6 November 2014.
- Mezali K. and Thandar A.S. 2014. First record of *Holothuria* (*Roweothuria*) *arguinensis* (Echinodermata: Holothuroidea: Aspidochirotida: Holothuriidae) from the Algerian coastal waters. *Marine Biology* 7, e40:1–4.
- Mezali K., Soualili D.L., Neghli L. and Conand C. 2014. Reproductive cycle of the sea cucumber *Holothuria* (*Platyperona*) *sanctori* (Holothuroidea: Echinodermata) in the southwestern Mediterranean Sea: Interpopulation variability. *Invertebrate Reproduction and Development* 58(3):179–189.
- Neghli L. and Mezali K. 2019. Algeria's sea cucumber fishery: Challenges for a new fishery. *SPC Beche-de-mer Information Bulletin* 39:71–75.
- Neghli L., Mezali K. and Soualili D.L. 2013. Contribution à l'étude de la reproduction de quelques espèces d'holothuries aspidochirotes inféodées aux herbiers de Posidonies de la côte algéroise. Poster. Workshop international sur la Biodiversité et Ecosystèmes Littoraux, 26-27-28 Novembre 2013, Oran (Algérie).
- Purcell S.W., Williamson D.H. and Ngaluaf P. 2018. Chinese market prices of beche-de-mer: Implications for fisheries and aquaculture. *Marine Policy* 91:58–56.
- Slimane-Tamacha F., Soualili D.L. and Mezali K. 2019. Reproductive biology of *Holothuria* (*Roweothuria*) *poli* (Holothuroidea: Echinodermata) from Oran Bay, Algeria. *SPC Beche-de-mer Information Bulletin* 39:47–53.