

RESEARCH ARTICLE

On the Hermite-Hadamard-Fejer-type inequalities for co-ordinated convex functions via fractional integrals

Hatice Yaldız^{a*}, Mehmet Zeki Sarıkaya^a and Zoubir Dahmani^b

^aDepartment of Mathematics, Faculty of Science and Arts, Düzce University, Düzce, Turkey

^bLaboratory of Pure and Applied Mathematics, UMAB, University of Mostaganem, Algeria
 yaldizhatice@gmail.com, sarikayamz@gmail.com, zzdahmani@yahoo.fr

ARTICLE INFO

Article History:
 Received 20 January 2017
 Accepted 06 June 2017
 Available 17 July 2017

Keywords:

Convex function
 Co-ordinated convex mapping
 Hermite-Hadamard-Fejer inequality
 Riemann-Liouville fractional integrals

AMS Classification 2010:
 26A33, 26A51, 26D15

ABSTRACT

In this paper, using Riemann-Liouville integral operators, we establish new fractional integral inequalities of Hermite-Hadamard-Fejer type for co-ordinated convex functions on a rectangle of \mathbb{R}^2 . The results presented here would provide extensions of those given in earlier works.



1. Introduction

Let $\Phi : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a convex mapping defined on the interval I of real numbers and $a, b \in I$, with $a < b$. The following double inequality is well known in the literature as the Hermite-Hadamard inequality [13]:

$$\begin{aligned} & \Phi\left(\frac{a+b}{2}\right) \\ & \leq \frac{1}{b-a} \int_a^b \Phi(x) dx \\ & \leq \frac{\Phi(a) + \Phi(b)}{2}. \end{aligned} \tag{1}$$

The most well-known inequalities related to the integral mean of a convex function are the Hermite-Hadamard inequalities or its weighted versions, the so-called Hermite-Hadamard-Fejér inequalities (see, [14], [19], [21]). In [11], Fejer gave a weighted generalization of the inequalities (1) as the following:

Theorem 1. Let $\Phi : [a, b] \rightarrow \mathbb{R}$ be a convex function. Then the inequality hold:

$$\begin{aligned} & \Phi\left(\frac{a+b}{2}\right) \int_a^b \Psi(x) dx \\ & \leq \int_a^b \Phi(x) \Psi(x) dx \\ & \leq \frac{\Phi(a) + \Phi(b)}{2} \int_a^b \Psi(x) dx, \end{aligned}$$

where $\Psi : [a, b] \rightarrow \mathbb{R}$ is nonnegative, integrable and symmetric to $\frac{(a+b)}{2}$.

In the following, we will give some necessary definitions and mathematical preliminaries of fractional calculus theory which are used further in this paper. More details, one can consult [12, 18].

Definition 1. ([4, 12, 18]) Let $\Phi \in L_1([a, b])$. The Riemann-Liouville integrals $J_{a+}^\alpha \Phi$ and $J_{b-}^\alpha \Phi$ of order $\alpha > 0$ with $a \geq 0$ are defined by

*Corresponding Author

$$J_{a+}^{\alpha} \Phi(x) = \frac{1}{\Gamma(\alpha)} \int_a^x (x-t)^{\alpha-1} \Phi(t) dt, \quad x > a$$

and

$$J_{b-}^{\alpha} \Phi(x) = \frac{1}{\Gamma(\alpha)} \int_x^b (t-x)^{\alpha-1} \Phi(t) dt, \quad x < b$$

respectively. Here, $\Gamma(\alpha)$ is the Gamma function and $J_{a+}^0 \Phi(x) = J_{b-}^0 \Phi(x) = \Phi(x)$.

Meanwhile, in [24], Sarıkaya et al. gave the following interesting Riemann-Liouville integral inequalities of Hermite-Hadamard-type:

Theorem 2. Let $K : [a, b] \rightarrow \mathbb{R}$ be a positive function with $0 \leq a < b$ and $K \in L_1([a, b])$. If K is a convex function on $[a, b]$, then the following inequalities for fractional integrals hold:

$$\begin{aligned} & K\left(\frac{a+b}{2}\right) \tag{2} \\ & \leq \frac{\Gamma(\alpha+1)}{2(b-a)^{\alpha}} [J_{a+}^{\alpha} K(b) + J_{b-}^{\alpha} K(a)] \\ & \leq \frac{K(a) + K(b)}{2} \end{aligned}$$

with $\alpha > 0$.

Later, in [14], Iscan presented the following Hermite-Hadamard-Fejer type inequalities for convex functions via Riemann-Liouville fractional integrals:

Theorem 3. Let $K : [a, b] \rightarrow \mathbb{R}$ be convex function with $0 \leq a < b$ and $K \in (L_1[a, b])$. If $L : [a, b] \rightarrow \mathbb{R}$ is nonnegative, integrable and symmetric with respect to $\frac{a+b}{2}$, then the following inequalities for fractional integrals hold:

$$\begin{aligned} & K\left(\frac{a+b}{2}\right) [J_{a+}^{\alpha} L(b) + J_{b-}^{\alpha} L(a)] \tag{3} \\ & \leq [J_{a+}^{\alpha} K(b) + J_{b-}^{\alpha} K(a)] \\ & \leq \frac{K(a) + K(b)}{2} [J_{a+}^{\alpha} L(b) + J_{b-}^{\alpha} L(a)] \end{aligned}$$

with $\alpha > 0$.

Let us now consider a bi-dimensional interval which will be used throughout this paper. So, we define $\Delta =: [a, b] \times [c, d]$ in \mathbb{R}^2 with $a < b$ and $c < d$. A mapping $\Phi : \Delta \rightarrow \mathbb{R}$ is said to be convex on the co-ordinates Δ if the following inequality:

$$\Phi(tx + (1-t)z, ty + (1-t)r)$$

$$\leq t\Phi(x, y) + (1-t)\Phi(z, r)$$

holds, for all $(x, y), (z, r) \in \Delta$ and $t \in [0, 1]$.

A function $\Phi : \Delta \rightarrow \mathbb{R}$ is said to be convex on the co-ordinates on Δ if the partial mappings $\Phi_y : [a, b] \rightarrow \mathbb{R}$, $\Phi_y(u) = \Phi(u, y)$ and $\Phi_x : [c, d] \rightarrow \mathbb{R}$, $\Phi_x(v) = \Phi(x, v)$ are convex where defined for all $x \in [a, b]$ and $y \in [c, d]$ (see, [10]).

A formal definition for co-ordinated convex functions may be stated as follows:

Definition 2. ([10]) A function $\Phi : \Delta \rightarrow \mathbb{R}$ will be called co-ordinated convex on Δ , for all $t, s \in [0, 1]$ and $(x, y), (u, r) \in \Delta$, if the following inequality holds:

$$\begin{aligned} & \Phi(tx + (1-t)y, su + (1-s)r) \\ & \leq ts\Phi(x, u) + s(1-t)\Phi(y, u) \tag{4} \end{aligned}$$

$$+t(1-s)\Phi(x, r) + (1-t)(1-s)\Phi(y, r).$$

Clearly, every convex function is a co-ordinated convex. Furthermore, there exists a co-ordinated convex function which is not convex, (see, [10]).

For several recent results concerning Hermite-Hadamard's inequality for some convex function on the co-ordinates on a rectangle of \mathbb{R}^2 , we refer the reader to ([1]- [3], [10], [15]- [17], [20], [22], [27]).

In [10], Dragomir established the following inequality of Hermite-Hadamard-type for co-ordinated convex mapping on a rectangle of \mathbb{R}^2 similar to (1).

Theorem 4. Suppose that $\Phi : \Delta \rightarrow \mathbb{R}$ is co-ordinated convex on Δ . Then one has the inequalities:

$$\begin{aligned}
 & \Phi\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \\
 & \leq \frac{1}{2} \left[\frac{1}{b-a} \int_a^b \Phi\left(x, \frac{c+d}{2}\right) dx \right. \\
 & \quad \left. + \frac{1}{d-c} \int_c^d \Phi\left(\frac{a+b}{2}, y\right) dy \right] \\
 & \leq \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d \Phi(x, y) dy dx \quad (5) \\
 & \leq \frac{1}{4} \left[\frac{1}{b-a} \int_a^b \Phi(x, c) dx + \frac{1}{b-a} \int_a^b \Phi(x, d) dx \right. \\
 & \quad \left. + \frac{1}{d-c} \int_c^d \Phi(a, y) dy + \frac{1}{d-c} \int_c^d \Phi(b, y) dy \right] \\
 & \leq \frac{\Phi(a, c) + \Phi(a, d) + \Phi(b, c) + \Phi(b, d)}{4}.
 \end{aligned}$$

The above inequalities are sharp.

Later, in [27], Sarikaya and Yaldiz proved inequalities of the Hermite-Hadamard type by using the definition of co-ordinated convex functions for L-Lipschitzian mappings.

In [3], a Hermite-Hadamard-Fejer type inequality for co-ordinated convex mappings is established as follows:

Theorem 5. *Let $\Phi : \Delta \rightarrow R$ be a co-ordinated convex function. Then the following double inequality hold:*

$$\begin{aligned}
 & \Phi\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \\
 & \leq \frac{\int_a^b \int_c^d \Phi(x, y) p(x, y) dy dx}{\int_a^b \int_c^d p(x, y) dy dx} \quad (6) \\
 & \leq \frac{\Phi(a, c) + \Phi(a, d) + \Phi(b, c) + \Phi(b, d)}{4},
 \end{aligned}$$

where $p : \Delta \rightarrow R$ is positive, integrable and symmetric with respect to $x = \frac{a+b}{2}$ and $y = \frac{c+d}{2}$ on the co-ordinates on Δ . The above inequalities are sharp.

Because of the wide application of Hermite Hadamard type inequalities, Fejer type inequalities and Riemann-Liouville integrals for two-variable functions, many authors extend their studies to Hermite Hadamard type inequalities and Fejer type inequalities involving Riemann-Liouville integrals not limited to integer integrals.

Definition 3. ([12, 18]) Let $\Phi \in L_1(\Delta)$. The Riemann-Liouville integrals $J_{a+,c+}^{\alpha,\beta}$, $J_{a+,d-}^{\alpha,\beta}$, $J_{b-,c+}^{\alpha,\beta}$ and $J_{b-,d-}^{\alpha,\beta}$ of order $\alpha, \beta > 0$ with $a, c \geq 0$ are defined by

$$\begin{aligned}
 J_{a+,c+}^{\alpha,\beta} \Phi(x, y) &= \\
 & \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_a^x \int_c^y (x-t)^{\alpha-1} (y-s)^{\beta-1} \Phi(t, s) ds dt,
 \end{aligned}$$

$$x > a, y > c$$

$$\begin{aligned}
 J_{a+,d-}^{\alpha,\beta} \Phi(x, y) &= \\
 & \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_a^x \int_y^d (x-t)^{\alpha-1} (s-y)^{\beta-1} \Phi(t, s) ds dt,
 \end{aligned}$$

$$x > a, y < d$$

$$\begin{aligned}
 J_{b-,c+}^{\alpha,\beta} \Phi(x, y) &= \\
 & \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_x^b \int_c^y (t-x)^{\alpha-1} (y-s)^{\beta-1} \Phi(t, s) ds dt,
 \end{aligned}$$

$$x < b, y > c$$

and

$$\begin{aligned}
 J_{b-,d-}^{\alpha,\beta} \Phi(x, y) &= \\
 & \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_x^b \int_y^d (t-x)^{\alpha-1} (s-y)^{\beta-1} \Phi(t, s) ds dt,
 \end{aligned}$$

$$x < b, y < d$$

respectively. Here, Γ is the Gamma function,

$$\begin{aligned}
 J_{a+,c+}^{0,0} \Phi(x, y) &= J_{a+,d-}^{0,0} \Phi(x, y) \\
 &= J_{b-,c+}^{0,0} \Phi(x, y) = J_{b-,d-}^{0,0} \Phi(x, y) = \Phi(x, y)
 \end{aligned}$$

and

$$J_{a+,c+}^{1,1} \Phi(x, y) = \int_a^x \int_c^y \Phi(t, s) ds dt.$$

Similar to Definition 1 and Definition 3, we introduce the following fractional integrals:

$$\begin{aligned}
 & J_{a+}^{\alpha} \Phi\left(x, \frac{c+d}{2}\right) \\
 &= \frac{1}{\Gamma(\alpha)} \int_a^x (x-t)^{\alpha-1} \Phi\left(t, \frac{c+d}{2}\right) dt, \quad x > a;
 \end{aligned}$$

$$\begin{aligned}
& J_{b-}^{\alpha} \Phi \left(x, \frac{c+d}{2} \right) \\
&= \frac{1}{\Gamma(\alpha)} \int_x^b (t-x)^{\alpha-1} \Phi \left(t, \frac{c+d}{2} \right) dt, \quad x < b; \\
& J_{c+}^{\beta} \Phi \left(\frac{a+b}{2}, y \right) \\
&= \frac{1}{\Gamma(\beta)} \int_c^y (y-s)^{\beta-1} \Phi \left(\frac{a+b}{2}, s \right) ds, \quad y > c; \\
& J_{d-}^{\beta} \Phi \left(\frac{a+b}{2}, y \right) \\
&= \frac{1}{\Gamma(\beta)} \int_y^d (s-y)^{\beta-1} \Phi \left(\frac{a+b}{2}, s \right) ds, \quad y < d.
\end{aligned}$$

It is remarkable that Sarıkaya et al. ([26]) and ([28]) gave the following interesting integral inequalities of Hermite-Hadamard-type involving Riemann-Liouville fractional integrals by using convex functions of 2-variables on the co-ordinates.

Theorem 6. Let $\Phi : \Delta \rightarrow \mathbb{R}$ be co-ordinated convex on Δ , with $0 \leq a < b$, $0 \leq c < d$ and $\Phi \in L_1(\Delta)$. Then one has the inequalities:

$$\begin{aligned}
& \Phi \left(\frac{a+b}{2}, \frac{c+d}{2} \right) \tag{7} \\
& \leq \frac{\Gamma(\alpha+1)\Gamma(\beta+1)}{4(b-a)^{\alpha}(d-c)^{\beta}} \\
& \times \left[J_{a+,c+}^{\alpha,\beta} \Phi(b,d) + J_{a+,d-}^{\alpha,\beta} \Phi(b,c) \right. \\
& \left. + J_{b-,c+}^{\alpha,\beta} \Phi(a,d) + J_{b-,d-}^{\alpha,\beta} \Phi(a,c) \right] \\
& \leq \frac{\Phi(a,c) + \Phi(a,d) + \Phi(b,c) + \Phi(b,d)}{4}.
\end{aligned}$$

For some recent results connected with fractional integral inequalities, see ([5]- [9], [23]- [26]).

The main aim of this paper is to establish new results on Hermite-Hadamard-Fejer type inequalities for co-ordinated convex functions on the rectangle Δ introduced in the first section of this paper. We will use the Riemann-Liouville integral operators to prove our main results.

2. Hermite-Hadamard-Fejer type inequalities for fractional integrals

In this section, using Riemann-Liouville fractional integral operators, we establish new results on Hermite-Hadamard-Fejer type inequalities for co-ordinated convex functions. We present evidence

by using two different methods. We begin by the following theorem:

Theorem 7. Let $\Phi : \Delta \rightarrow \mathbb{R}$ be a co-ordinated convex function such that $\Phi \in L_1(\Delta)$. If $\Psi : \Delta \rightarrow \mathbb{R}$ is nonnegative, integrable and symmetric with respect to $\frac{a+b}{2}, \frac{c+d}{2}$ on the co-ordinates, then for any $\alpha, \beta > 0$ with $a, c \geq 0$, the following integral inequalities hold

$$\begin{aligned}
& \Phi \left(\frac{a+b}{2}, \frac{c+d}{2} \right) \tag{8} \\
& \times \left[J_{a+,c+}^{\alpha,\beta} \Psi(b,d) + J_{a+,d-}^{\alpha,\beta} \Psi(b,c) \right. \\
& \left. + J_{b-,c+}^{\alpha,\beta} \Psi(a,d) + J_{b-,d-}^{\alpha,\beta} \Psi(a,c) \right] \\
& \leq \frac{1}{4} \left[J_{a+,c+}^{\alpha,\beta} (\Phi\Psi)(b,d) + J_{a+,d-}^{\alpha,\beta} (\Phi\Psi)(b,c) \right. \\
& \left. + J_{b-,c+}^{\alpha,\beta} (\Phi\Psi)(a,d) + J_{b-,d-}^{\alpha,\beta} (\Phi\Psi)(a,c) \right] \\
& \leq \frac{\Phi(a,c) + \Phi(a,d) + \Phi(b,c) + \Phi(b,d)}{4} \\
& \times \left[J_{a+,c+}^{\alpha,\beta} \Psi(b,d) + J_{a+,d-}^{\alpha,\beta} \Psi(b,c) \right. \\
& \left. + J_{b-,c+}^{\alpha,\beta} \Psi(a,d) + J_{b-,d-}^{\alpha,\beta} \Psi(a,c) \right].
\end{aligned}$$

Proof. Since Φ is a convex function on Δ , then, for all $(t, s) \in [0, 1] \times [0, 1]$, we can write:

$$\begin{aligned}
& \Phi \left(\frac{a+b}{2}, \frac{c+d}{2} \right) \tag{9} \\
&= \Phi \left(\frac{ta + (1-t)b + (1-t)a + tb}{2}, \right. \\
& \left. \frac{sc + (1-s)d + (1-s)c + sd}{2} \right) \\
& \leq \frac{1}{4} \left[\Phi(ta + (1-t)b, sc + (1-s)d) \right. \\
& \left. + \Phi(ta + (1-t)b, (1-s)c + sd) \right. \\
& \left. + \Phi((1-t)a + tb, sc + (1-s)d) \right. \\
& \left. + \Phi((1-t)a + tb, (1-s)c + sd) \right].
\end{aligned}$$

Multiplying both sides of (9) by $t^{\alpha-1}s^{\beta-1}\Psi((1-t)a + tb, (1-s)c + sd)$, and integrating the resulting inequality with respect to (t, s) on $[0, 1] \times [0, 1]$, we obtain

$$\begin{aligned}
 & \Phi\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \\
 & \times \int_0^1 \int_0^1 t^{\alpha-1} s^{\beta-1} \Psi((1-t)a+tb, (1-s)c+sd) ds dt \\
 & \leq \frac{1}{4} \int_0^1 \int_0^1 t^{\alpha-1} s^{\beta-1} [\Phi(ta+(1-t)b, sc+(1-s)d) \\
 & \quad + \Phi(ta+(1-t)b, (1-s)c+sd) \\
 & \quad + \Phi((1-t)a+tb, sc+(1-s)d) \\
 & \quad + \Phi((1-t)a+tb, (1-s)c+sd)] \\
 & \quad \times \Psi((1-t)a+tb, (1-s)c+sd) ds dt \\
 & = \frac{1}{4} \int_0^1 \int_0^1 t^{\alpha-1} s^{\beta-1} \Phi(ta+(1-t)b, sc+(1-s)d) \\
 & \quad \times \Psi((1-t)a+tb, (1-s)c+sd) ds dt \\
 & \quad + \int_0^1 \int_0^1 t^{\alpha-1} s^{\beta-1} \Phi(ta+(1-t)b, (1-s)c+sd) \\
 & \quad \times \Psi((1-t)a+tb, (1-s)c+sd) ds dt \\
 & \quad + \int_0^1 \int_0^1 t^{\alpha-1} s^{\beta-1} \Phi((1-t)a+tb, sc+(1-s)d) \\
 & \quad \times \Psi((1-t)a+tb, (1-s)c+sd) ds dt \\
 & \quad + \int_0^1 \int_0^1 t^{\alpha-1} s^{\beta-1} \Phi((1-t)a+tb, (1-s)c+sd) \\
 & \quad \times \Psi((1-t)a+tb, (1-s)c+sd) ds dt. \\
 & \frac{1}{(b-a)^\alpha (d-c)^\beta} \Phi\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \\
 & \times \int_a^b \int_c^d (x-a)^{\alpha-1} (y-c)^{\beta-1} \Psi(x, y) dy dx \\
 & \leq \frac{1}{4(b-a)^\alpha (d-c)^\beta} \\
 & \times \left\{ \int_a^b \int_c^d (x-a)^{\alpha-1} (y-c)^{\beta-1} \right. \\
 & \quad \times \Phi(a+b-x, c+d-y) \Psi(x, y) dy dx \\
 & \quad + \int_a^b \int_c^d (x-a)^{\alpha-1} (d-y)^{\beta-1} \\
 & \quad \times \Phi(a+b-x, y) \Psi(x, y) dy dx \\
 & \quad + \int_a^b \int_c^d (b-x)^{\alpha-1} (y-c)^{\beta-1} \\
 & \quad \times \Phi(x, c+d-y) \Psi(x, y) dy dx \\
 & \quad \left. + \int_a^b \int_c^d (b-x)^{\alpha-1} (d-y)^{\beta-1} \Phi(x, y) \Psi(x, y) dy dx \right\} \\
 & = \frac{1}{4(b-a)^\alpha (d-c)^\beta} \\
 & \times \left\{ \int_a^b \int_c^d (b-x)^{\alpha-1} (y-c)^{\beta-1} \right. \\
 & \quad \times \Phi(x, c+d-y) \Psi(a+b-x, y) dy dx \\
 & \quad + \int_a^b \int_c^d (b-x)^{\alpha-1} (d-y)^{\beta-1} \\
 & \quad \times \Phi(x, y) \Psi(a+b-x, y) dy dx \\
 & \quad + \int_a^b \int_c^d (b-x)^{\alpha-1} (y-c)^{\beta-1} \\
 & \quad \times \Phi(x, c+d-y) \Psi(x, y) dy dx \\
 & \quad \left. + \int_a^b \int_c^d (b-x)^{\alpha-1} (d-y)^{\beta-1} \Phi(x, y) \Psi(x, y) dy dx \right\}
 \end{aligned}$$

Setting $x = tb + (1-t)a$, $y = sd + (1-s)c$ and $dx = (b-a) dt$, $dy = (d-c) ds$, we obtain:

$$\begin{aligned}
&= \frac{1}{4(b-a)^\alpha (d-c)^\beta} \left\{ \int_a^b \int_c^d (b-x)^{\alpha-1} (y-c)^{\beta-1} \right. \\
&\quad \times \Phi(x, c+d-y) \Psi(x, y) dy dx \\
&\quad + \int_a^b \int_c^d (b-x)^{\alpha-1} (d-y)^{\beta-1} \Phi(x, y) \Psi(x, y) dy dx \\
&\quad + \int_a^b \int_c^d (b-x)^{\alpha-1} (y-c)^{\beta-1} \\
&\quad \times \Phi(x, c+d-y) \Psi(x, y) dy dx \\
&\quad \left. + \int_a^b \int_c^d (b-x)^{\alpha-1} (d-y)^{\beta-1} \Phi(x, y) \Psi(x, y) dy dx \right\}. \\
&\int_0^1 \int_0^1 t^{\alpha-1} s^{\beta-1} [\Phi(ta + (1-t)b, sc + (1-s)d) \\
&\quad + \Phi(ta + (1-t)b, (1-s)c + sd) \\
&\quad + \Phi((1-t)a + tb, sc + (1-s)d) \\
&\quad + \Phi((1-t)a + tb, (1-s)c + sd)] \\
&\quad \times \Psi(tb + (1-t)a, sd + (1-s)c) ds dt \\
&\leq [\Phi(a, c) + \Phi(b, c) + \Phi(a, d) + \Phi(b, d)] \\
&\quad \times \int_0^1 \int_0^1 t^{\alpha-1} s^{\beta-1} \Psi(tb + (1-t)a, sd + (1-s)c) ds dt.
\end{aligned}$$

Therefore,

$$\begin{aligned}
&\frac{\Gamma(\alpha)\Gamma(\beta)}{(b-a)^\alpha (d-c)^\beta} \Phi\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \\
&\quad \times \left[J_{a^+c^+}^{\alpha,\beta} \Psi(b, d) + J_{a^+d^-}^{\alpha,\beta} \Psi(b, c) \right. \\
&\quad \left. + J_{b^-c^+}^{\alpha,\beta} \Psi(a, d) + J_{b^-d^-}^{\alpha,\beta} \Psi(a, c) \right] \\
&\leq \frac{\Gamma(\alpha)\Gamma(\beta)}{4(b-a)^\alpha (d-c)^\beta} \\
&\quad \times \left[J_{a^+c^+}^{\alpha,\beta} (\Phi\Psi)(b, d) + J_{a^+d^-}^{\alpha,\beta} (\Phi\Psi)(b, c) \right. \\
&\quad \left. + J_{b^-c^+}^{\alpha,\beta} (\Phi\Psi)(a, d) + J_{b^-d^-}^{\alpha,\beta} (\Phi\Psi)(a, c) \right].
\end{aligned}$$

The first inequality of (8) is thus proved.

We shall prove the second inequality of (8): Since f is a convex function on Δ , then, for all $(t, s) \in [0, 1] \times [0, 1]$, it yields

$$\begin{aligned}
&\Phi(ta + (1-t)b, sc + (1-s)d) \\
&\quad + \Phi(ta + (1-t)b, (1-s)c + sd) \\
&\quad + \Phi((1-t)a + tb, sc + (1-s)d) \\
&\quad + \Phi((1-t)a + tb, (1-s)c + sd) \quad (10) \\
&\leq \Phi(a, c) + \Phi(b, c) + \Phi(a, d) + \Phi(b, d).
\end{aligned}$$

Then, multiplying both sides of (10) by $t^{\alpha-1}s^{\beta-1}\Psi(tb + (1-t)a, sd + (1-s)c)$ and integrating the resulting inequality with respect to (t, s) over $[0, 1] \times [0, 1]$, we get

That is,

$$\begin{aligned}
&\frac{1}{4} \left[J_{a^+c^+}^{\alpha,\beta} (\Phi\Psi)(b, d) + J_{a^+d^-}^{\alpha,\beta} (\Phi\Psi)(b, c) \right. \\
&\quad \left. + J_{b^-c^+}^{\alpha,\beta} (\Phi\Psi)(a, d) + J_{b^-d^-}^{\alpha,\beta} (\Phi\Psi)(a, c) \right] \\
&\leq \frac{\Phi(a, c) + \Phi(a, d) + \Phi(b, c) + \Phi(b, d)}{4} \\
&\quad \times \left[J_{a^+c^+}^{\alpha,\beta} \Psi(b, d) + J_{a^+d^-}^{\alpha,\beta} \Psi(b, c) \right. \\
&\quad \left. + J_{b^-c^+}^{\alpha,\beta} \Psi(a, d) + J_{b^-d^-}^{\alpha,\beta} \Psi(a, c) \right].
\end{aligned}$$

The proof of Theorem 7 is thus achieved. \square

Remark 1. In Theorem 7:

(i) If we take $\alpha = \beta = 1$, then the inequality (8) becomes the inequality (6) of Theorem 5.

(ii) If we take $\Psi(x, y) = 1$, then (8) becomes (7) of Theorem 6.

We prove also the following result:

Theorem 8. Let $\Phi : \Delta \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ be a co-ordinated convex function on Δ , with $a, c \geq 0$, $\alpha, \beta > 0$ and $\Phi \in L_1(\Delta)$. If $\Psi : \Delta \rightarrow \mathbb{R}$ is non-negative, integrable and symmetric with respect to $\frac{a+b}{2}$ and $\frac{c+d}{2}$ on the co-ordinates, then we have:

$$\begin{aligned}
&\Phi\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \times \left[J_{a^+c^+}^{\alpha,\beta} \Psi(b, d) + J_{a^+d^-}^{\alpha,\beta} \Psi(b, c) \right. \\
&\quad \left. + J_{b^-c^+}^{\alpha,\beta} \Psi(a, d) + J_{b^-d^-}^{\alpha,\beta} \Psi(a, c) \right] \quad (11)
\end{aligned}$$

$$\begin{aligned}
 &\leq J_{a+}^{\alpha} \left[\Phi \left(b, \frac{c+d}{2} \right) J_{c+}^{\beta} \Psi(b, d) \right] \\
 &\quad + J_{a+}^{\alpha} \left[\Phi \left(b, \frac{c+d}{2} \right) J_{d-}^{\beta} \Psi(b, c) \right] \\
 &\quad + J_{b-}^{\alpha} \left[\Phi \left(a, \frac{c+d}{2} \right) J_{c+}^{\beta} \Psi(a, d) \right] \\
 &\quad + J_{b-}^{\alpha} \left[\Phi \left(a, \frac{c+d}{2} \right) J_{d-}^{\beta} \Psi(a, c) \right] \\
 &\quad + J_{c+}^{\beta} \left[\Phi \left(\frac{a+b}{2}, d \right) J_{a+}^{\alpha} \Psi(b, d) \right] \\
 &\quad + J_{c+}^{\beta} \left[\Phi \left(\frac{a+b}{2}, d \right) J_{b-}^{\alpha} \Psi(a, d) \right] \\
 &\quad + J_{d-}^{\beta} \left[\Phi \left(\frac{a+b}{2}, c \right) J_{a+}^{\alpha} \Psi(b, c) \right] \\
 &\quad + J_{d-}^{\beta} \left[\Phi \left(\frac{a+b}{2}, c \right) J_{b-}^{\alpha} \Psi(a, c) \right] \\
 &\leq 2 \left[J_{a+,c+}^{\alpha,\beta} (\Phi\Psi) (b, d) + J_{a+,d-}^{\alpha,\beta} (\Phi\Psi) (b, c) \right. \\
 &\quad \left. + J_{b-,c+}^{\alpha,\beta} (\Phi\Psi) (a, d) + J_{b-,d-}^{\alpha,\beta} (\Phi\Psi) (a, c) \right] \\
 &\leq J_{a+}^{\alpha} \left[\Phi (b, c) J_{c+}^{\beta} \Psi(b, d) \right] \\
 &\quad + J_{a+}^{\alpha} \left[\Phi (b, d) J_{d-}^{\beta} \Psi(b, c) \right] \\
 &\quad + J_{b-}^{\alpha} \left[\Phi (a, c) J_{c+}^{\beta} \Psi(a, d) \right] \\
 &\quad + J_{b-}^{\alpha} \left[\Phi (a, d) J_{d-}^{\beta} \Psi(a, c) \right] \\
 &\quad + J_{c+}^{\beta} \left[\Phi (a, d) J_{a+}^{\alpha} \Psi(b, d) \right] \\
 &\quad + J_{c+}^{\beta} \left[\Phi (b, d) J_{b-}^{\alpha} \Psi(a, d) \right] \\
 &\quad + J_{d-}^{\beta} \left[\Phi (a, c) J_{a+}^{\alpha} \Psi(b, c) \right] \\
 &\quad + J_{d-}^{\beta} \left[\Phi (b, c) J_{b-}^{\alpha} \Psi(a, c) \right] \\
 &\leq \frac{\Phi (a, c) + \Phi (a, d) + \Phi (b, c) + \Phi (b, d)}{4} \\
 &\quad \times \left[J_{a+,c+}^{\alpha,\beta} \Psi(b, d) + J_{a+,d-}^{\alpha,\beta} \Psi(b, c) \right. \\
 &\quad \left. + J_{b-,c+}^{\alpha,\beta} \Psi(a, d) + J_{b-,d-}^{\alpha,\beta} \Psi(a, c) \right].
 \end{aligned}$$

Proof. Since $\Phi : \Delta \rightarrow \mathbb{R}$ is convex on the co-ordinates, it follows that the mapping $F_x :$

$[c, d] \rightarrow \mathbb{R}$, $F_x(y) = \Phi(x, y)$, is convex on $[c, d]$ and the mapping $G_x : [c, d] \rightarrow \mathbb{R}$, $G_x(y) = \Psi(x, y)$ is nonnegative, integrable and symmetric with respect to $\frac{c+d}{2}$, for all $x \in [a, b]$. Then, thanks to the inequalities (3), we can write

$$\begin{aligned}
 &F_x \left(\frac{c+d}{2} \right) \left[J_{c+}^{\beta} G_x(d) + J_{d-}^{\beta} G_x(c) \right] \\
 &\leq J_{c+}^{\beta} (F_x G_x) (d) + J_{d-}^{\beta} (F_x G_x) (c) \\
 &\leq \frac{F_x (c) + F_x (d)}{2} \left[J_{c+}^{\beta} G_x(d) + J_{d-}^{\beta} G_x(c) \right], \\
 &x \in [a, b].
 \end{aligned}$$

That is,

$$\begin{aligned}
 &\Phi \left(x, \frac{c+d}{2} \right) \frac{1}{\Gamma(\beta)} \left[\int_c^d (d-y)^{\beta-1} \Psi(x, y) dy \right. \\
 &\quad \left. + \int_c^d (y-c)^{\beta-1} \Psi(x, y) dy \right] \\
 &\leq \frac{1}{\Gamma(\beta)} \left[\int_c^d (d-y)^{\beta-1} \Psi(x, y) \Phi(x, y) dy \right. \\
 &\quad \left. + \int_c^d (y-c)^{\beta-1} \Psi(x, y) \Phi(x, y) dy \right] \\
 &\leq \frac{\Phi(x, c) + \Phi(x, d)}{2} \frac{1}{\Gamma(\beta)} \left[\int_c^d (d-y)^{\beta-1} \Psi(x, y) dy \right. \\
 &\quad \left. + \int_c^d (y-c)^{\beta-1} \Psi(x, y) dy \right],
 \end{aligned}$$

for all $x \in [a, b]$.

Multiplying both sides of (12) by $\frac{(b-x)^{\alpha-1}}{\Gamma(\alpha)}$ and $\frac{(x-a)^{\alpha-1}}{\Gamma(\alpha)}$, and integrating with respect to x over $[a, b]$, respectively, we have

$$\begin{aligned}
 &\frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_a^b \int_c^d (b-x)^{\alpha-1} (d-y)^{\beta-1} \\
 &\quad \times \Phi \left(x, \frac{c+d}{2} \right) \Psi(x, y) dy dx \tag{13} \\
 &\quad + \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_a^b \int_c^d (b-x)^{\alpha-1} (y-c)^{\beta-1} \\
 &\quad \times \Phi \left(x, \frac{c+d}{2} \right) \Psi(x, y) dy dx
 \end{aligned}$$

$$\leq \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \left[\int_a^b \int_c^d (b-x)^{\alpha-1}(d-y)^{\beta-1} \times \Phi(x,y) \Psi(x,y) dydx \right. \\ \left. + \int_a^b \int_c^d (b-x)^{\alpha-1}(y-c)^{\beta-1} \Phi(x,y) \Psi(x,y) dydx \right] \tag{14}$$

$$\leq \frac{1}{2\Gamma(\alpha)\Gamma(\beta)} \left[\int_a^b \int_c^d (b-x)^{\alpha-1}(d-y)^{\beta-1} \times \Phi(x,c) \Psi(x,y) dydx \right. \\ \left. + \int_a^b \int_c^d (b-x)^{\alpha-1}(y-c)^{\beta-1} \Phi(x,c) \Psi(x,y) dydx \right. \\ \left. + \int_a^b \int_c^d (b-x)^{\alpha-1}(d-y)^{\beta-1} \Phi(x,d) \Psi(x,y) dydx \right. \\ \left. + \int_a^b \int_c^d (b-x)^{\alpha-1}(y-c)^{\beta-1} \Phi(x,d) \Psi(x,y) dydx \right].$$

and

$$\frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_a^b \int_c^d (x-a)^{\alpha-1}(d-y)^{\beta-1} \times \Phi\left(x, \frac{c+d}{2}\right) \Psi(x,y) dydx \tag{15} \\ + \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_a^b \int_c^d (x-a)^{\alpha-1}(y-c)^{\beta-1} \times \Phi\left(x, \frac{c+d}{2}\right) \Psi(x,y) dydx \\ \leq \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \times \left[\int_a^b \int_c^d (x-a)^{\alpha-1}(d-y)^{\beta-1} \times \Phi(x,y) \Psi(x,y) dydx \right. \\ \left. + \int_a^b \int_c^d (x-a)^{\alpha-1}(y-c)^{\beta-1} \times \Phi(x,y) \Psi(x,y) dydx \right] \\ \leq \frac{1}{2\Gamma(\alpha)\Gamma(\beta)} \left[\int_a^b \int_c^d (x-a)^{\alpha-1}(d-y)^{\beta-1} \times \Phi(x,c) \Psi(x,y) dydx \right. \\ \left. + \int_a^b \int_c^d (x-a)^{\alpha-1}(y-c)^{\beta-1} \Phi(x,c) \Psi(x,y) dydx \right. \\ \left. + \int_a^b \int_c^d (x-a)^{\alpha-1}(d-y)^{\beta-1} \Phi(x,d) \Psi(x,y) dydx \right. \\ \left. + \int_a^b \int_c^d (x-a)^{\alpha-1}(y-c)^{\beta-1} \Phi(x,d) \Psi(x,y) dydx \right].$$

For the mappings $F_y : [a, b] \rightarrow \mathbb{R}$, $F_y(x) = \Phi(x, y)$ and $G_y : [a, b] \rightarrow \mathbb{R}$, $G_y(x) = \Psi(x, y)$, we use the same arguments as before. So, we can state that

$$\frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_a^b \int_c^d (b-x)^{\alpha-1}(d-y)^{\beta-1} \times \Phi\left(\frac{a+b}{2}, y\right) \Psi(x,y) dydx \tag{16}$$

$$+ \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_a^b \int_c^d (x-a)^{\alpha-1}(d-y)^{\beta-1} \times \Phi\left(\frac{a+b}{2}, y\right) \Psi(x,y) dydx \\ \leq \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \left[\int_a^b \int_c^d (b-x)^{\alpha-1}(d-y)^{\beta-1} \times \Phi(x,y) \Psi(x,y) dydx \right. \\ \left. + \int_a^b \int_c^d (x-a)^{\alpha-1}(d-y)^{\beta-1} \Phi(x,y) \Psi(x,y) dydx \right]$$

$$\leq \frac{1}{2\Gamma(\alpha)\Gamma(\beta)} \left[\int_a^b \int_c^d (b-x)^{\alpha-1}(d-y)^{\beta-1} \times \Phi(a,y) \Psi(x,y) dydx \right. \\ \left. + \int_a^b \int_c^d (x-a)^{\alpha-1}(d-y)^{\beta-1} \Phi(a,y) \Psi(x,y) dydx \right. \\ \left. + \int_a^b \int_c^d (b-x)^{\alpha-1}(d-y)^{\beta-1} \Phi(b,y) \Psi(x,y) dydx \right. \\ \left. + \int_a^b \int_c^d (x-a)^{\alpha-1}(d-y)^{\beta-1} \Phi(b,y) \Psi(x,y) dydx \right]$$

and

$$\frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_a^b \int_c^d (b-x)^{\alpha-1}(y-c)^{\beta-1} \times \Phi\left(\frac{a+b}{2}, y\right) \Psi(x,y) dydx \tag{17}$$

$$+ \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_a^b \int_c^d (x-a)^{\alpha-1}(y-c)^{\beta-1} \times \Phi\left(\frac{a+b}{2}, y\right) \Psi(x,y) dydx \\ \leq \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \left[\int_a^b \int_c^d (b-x)^{\alpha-1}(y-c)^{\beta-1} \times \Phi(x,y) \Psi(x,y) dydx \right. \\ \left. + \int_a^b \int_c^d (x-a)^{\alpha-1}(y-c)^{\beta-1} \Phi(x,y) \Psi(x,y) dydx \right]$$

$$\begin{aligned} &\leq \frac{1}{2\Gamma(\alpha)\Gamma(\beta)} \left[\int_a^b \int_c^d (b-x)^{\alpha-1}(y-c)^{\beta-1} \right. \\ &\quad \times \Phi(a, y) \Psi(x, y) dydx \\ &\quad + \int_a^b \int_c^d (x-a)^{\alpha-1}(y-c)^{\beta-1} \Phi(a, y) \Psi(x, y) dydx \\ &\quad + \int_a^b \int_c^d (b-x)^{\alpha-1}(y-c)^{\beta-1} \Phi(b, y) \Psi(x, y) dydx \\ &\quad \left. + \int_a^b \int_c^d (x-a)^{\alpha-1}(y-c)^{\beta-1} \right. \\ &\quad \times \Phi(b, y) \Psi(x, y) dydx \Big] \\ &\quad + J_{c+}^\beta [\Phi(a, d) J_{a+}^\alpha \Psi(b, d)] \\ &\quad + J_{c+}^\beta [\Phi(b, d) J_{b-}^\alpha \Psi(a, d)] \\ &\quad + J_{d-}^\beta [\Phi(a, c) J_{a+}^\alpha \Psi(b, c)] \\ &\quad + J_{d-}^\beta [\Phi(b, c) J_{b-}^\alpha \Psi(a, c)]. \end{aligned}$$

Adding the inequalities (13)-(17), we can write

$$\begin{aligned} &J_{a+}^\alpha \left[\Phi \left(b, \frac{c+d}{2} \right) J_{c+}^\beta \Psi(b, d) \right] \\ &+ J_{a+}^\alpha \left[\Phi \left(b, \frac{c+d}{2} \right) J_{d-}^\beta \Psi(b, c) \right] \\ &+ J_{b-}^\alpha \left[\Phi \left(a, \frac{c+d}{2} \right) J_{c+}^\beta \Psi(a, d) \right] \\ &+ J_{b-}^\alpha \left[\Phi \left(a, \frac{c+d}{2} \right) J_{d-}^\beta \Psi(a, c) \right] \\ &+ J_{c+}^\beta \left[\Phi \left(\frac{a+b}{2}, d \right) J_{a+}^\alpha \Psi(b, d) \right] \\ &+ J_{c+}^\beta \left[\Phi \left(\frac{a+b}{2}, d \right) J_{b-}^\alpha \Psi(a, d) \right] \\ &+ J_{d-}^\beta \left[\Phi \left(\frac{a+b}{2}, c \right) J_{a+}^\alpha \Psi(b, c) \right] \\ &+ J_{d-}^\beta \left[\Phi \left(\frac{a+b}{2}, c \right) J_{b-}^\alpha \Psi(a, c) \right] \\ &\leq 2 \left[J_{a+,c+}^{\alpha,\beta} (\Phi\Psi)(b, d) + J_{a+,d-}^{\alpha,\beta} (\Phi\Psi)(b, c) \right. \\ &\quad \left. + J_{b-,c+}^{\alpha,\beta} (\Phi\Psi)(a, d) + J_{b-,d-}^{\alpha,\beta} (\Phi\Psi)(a, c) \right] \\ &\leq J_{a+}^\alpha \left[\Phi(b, c) J_{c+}^\beta \Psi(b, d) \right] \\ &\quad + J_{a+}^\alpha \left[\Phi(b, d) J_{d-}^\beta \Psi(b, c) \right] \\ &\quad + J_{b-}^\alpha \left[\Phi(a, c) J_{c+}^\beta \Psi(a, d) \right] \\ &\quad + J_{b-}^\alpha \left[\Phi(a, d) J_{d-}^\beta \Psi(a, c) \right] \end{aligned}$$

These give the second and the third inequalities in (11).

Now, by using the first inequality in (3), it yields that

$$\begin{aligned} &\Phi \left(\frac{a+b}{2}, \frac{c+d}{2} \right) \\ &\times \left[\int_a^b \int_c^d (b-x)^{\alpha-1}(d-y)^{\beta-1} \Psi(x, y) dydx \right. \\ &\quad \left. + \int_a^b \int_c^d (x-a)^{\alpha-1}(d-y)^{\beta-1} \Psi(x, y) dydx \right] \\ &\leq \int_a^b \int_c^d (b-x)^{\alpha-1}(d-y)^{\beta-1} \\ &\quad \times \Phi \left(x, \frac{c+d}{2} \right) \Psi(x, y) dydx \\ &\quad + \int_a^b \int_c^d (x-a)^{\alpha-1}(d-y)^{\beta-1} \Phi \left(x, \frac{c+d}{2} \right) \Psi(x, y) dydx \end{aligned}$$

and

$$\begin{aligned} &\Phi \left(\frac{a+b}{2}, \frac{c+d}{2} \right) \\ &\times \left[\int_a^b \int_c^d (b-x)^{\alpha-1}(d-y)^{\beta-1} \Psi(x, y) dydx \right. \\ &\quad \left. + \int_a^b \int_c^d (b-x)^{\alpha-1}(y-c)^{\beta-1} \Psi(x, y) dydx \right] \\ &\leq \int_a^b \int_c^d (b-x)^{\alpha-1}(d-y)^{\beta-1} \\ &\quad \times \Phi \left(\frac{a+b}{2}, y \right) \Psi(x, y) dydx \\ &\quad + \int_a^b \int_c^d (b-x)^{\alpha-1}(y-c)^{\beta-1} \\ &\quad \times \Phi \left(\frac{a+b}{2}, y \right) \Psi(x, y) dydx. \end{aligned}$$

By addition, and using the fact that Ψ is symmetric, we get

$$\begin{aligned}
& \Phi\left(\frac{a+b}{2}, \frac{c+d}{2}\right) && \leq \frac{\Phi(a,d) + \Phi(b,d)}{2}, \\
& \times \left[J_{a_+,c_+}^{\alpha,\beta} \Psi(b,d) + J_{a_+,d_-}^{\alpha,\beta} \Psi(b,c) \right. \\
& \quad \left. + J_{b_-,c_+}^{\alpha,\beta} \Psi(a,d) + J_{b_-,d_-}^{\alpha,\beta} \Psi(a,c) \right] \\
& \leq J_{a_+}^{\alpha} \left[\Phi\left(b, \frac{c+d}{2}\right) J_{c_+}^{\beta} \Psi(b,d) \right] \\
& \quad + J_{a_+}^{\alpha} \left[\Phi\left(b, \frac{c+d}{2}\right) J_{d_-}^{\beta} \Psi(b,c) \right] \\
& \quad + J_{b_-}^{\alpha} \left[\Phi\left(a, \frac{c+d}{2}\right) J_{c_+}^{\beta} \Psi(a,d) \right] \\
& \quad + J_{b_-}^{\alpha} \left[\Phi\left(a, \frac{c+d}{2}\right) J_{d_-}^{\beta} \Psi(a,c) \right] \\
& \quad + J_{c_+}^{\beta} \left[\Phi\left(\frac{a+b}{2}, d\right) J_{a_+}^{\alpha} \Psi(b,d) \right] \\
& \quad + J_{c_+}^{\beta} \left[\Phi\left(\frac{a+b}{2}, d\right) J_{b_-}^{\alpha} \Psi(a,d) \right] \\
& \quad + J_{d_-}^{\beta} \left[\Phi\left(\frac{a+b}{2}, c\right) J_{a_+}^{\alpha} \Psi(b,c) \right] \\
& \quad + J_{d_-}^{\beta} \left[\Phi\left(\frac{a+b}{2}, c\right) J_{b_-}^{\alpha} \Psi(a,c) \right] \\
& && \leq \frac{\Phi(a,d) + \Phi(b,d)}{2} \\
& && \frac{\beta}{2(d-c)^{\beta}} \left[\int_c^d (d-y)^{\beta-1} \Phi(a,y) dy \right. \\
& && \quad \left. + \int_c^d (y-c)^{\beta-1} \Phi(a,y) dy \right] \\
& && \leq \frac{\Phi(a,c) + \Phi(a,d)}{2}, \\
& && \frac{\beta}{2(d-c)^{\beta}} \left[\int_c^d (d-y)^{\beta-1} \Phi(b,y) dy \right. \\
& && \quad \left. + \int_c^d (y-c)^{\beta-1} \Phi(b,y) dy \right] \\
& && \leq \frac{\Phi(b,c) + \Phi(b,d)}{2}.
\end{aligned}$$

By addition, we get the last inequality in (11). \square

Remark 2. In Theorem 8, if we take $\alpha = \beta = 1$, then the inequalities (11) become (5).

3. Conclusion

In this paper, we established the Hermite-Hadamard-Fejer type inequalities for co-ordinated mappings related results to present new type of inequalities involving Riemann-Liouville integral operator. The results presented in this paper would provide generalizations of those given in earlier works. The findings of this study have several significant implications for future applications.

which gives the first inequality in (11).

Finally, by using the second inequality in (3), we can state that:

$$\begin{aligned}
& \frac{\alpha}{2(b-a)^{\alpha}} \left[\int_a^b (b-x)^{\alpha-1} \Phi(x,c) dx \right. \\
& \quad \left. + \int_a^b (x-a)^{\alpha-1} \Phi(x,c) dx \right] \\
& \leq \frac{\Phi(a,c) + \Phi(b,c)}{2}, \\
& \frac{\alpha}{2(b-a)^{\alpha}} \left[\int_a^b (b-x)^{\alpha-1} \Phi(x,d) dx \right. \\
& \quad \left. + \int_a^b (x-a)^{\alpha-1} \Phi(x,d) dx \right]
\end{aligned}$$

References

- [1] Alomari, M., Darus, M., Co-ordinated s-convex function in the first sense with some Hadamard-type inequalities, Int. J. Contemp. Math. Sciences, 3 (32),1557-1567 (2008).
- [2] Alomari, M., Darus, M., On the Hadamard's inequality for log-convex functions on the coordinates, J. of Inequal. and Appl, Article ID 283147, 13 pages,(2009).
- [3] Alomari, M., Darus, M., Fejer inequality for double integrals, Facta Universitatis (Nis) Ser. Math. Inform., 24 , 15-28(2009).
- [4] Kilbas, A.A., Srivastava H.M., Trujillo, J.J., Theory and Applications of Fractional Differential Equations, North-Holland Mathematics Studies, 204, Elsevier Sci. B.V., Amsterdam, (2006).
- [5] Belarbi, S., Dahmani, Z., On some new fractional integral inequalities, J. Ineq. Pure and Appl. Math., 10(3) , Art. 86, (2009).
- [6] Dahmani, Z., New inequalities in fractional integrals, International Journal of Nonlinear Science, 9(4) , 493-497, (2010).

- [7] Dahmani, Z., On Minkowski and Hermite-Hadamard integral inequalities via fractional integration, *Ann. Funct. Anal.* 1(1), 51-58, (2010).
- [8] Dahmani, Z., Tabharit, L., Taf, S., Some fractional integral inequalities, *Nonl. Sci. Lett. A*, 1(2), 155-160, (2010).
- [9] Dahmani, Z., Tabharit, L., Taf, S., New generalizations of Grüss inequality using Riemann-Liouville fractional integrals, *Bull. Math. Anal. Appl.*, 2(3), 93-99, (2010).
- [10] Dragomir, S.S., On Hadamard's inequality for convex functions on the co-ordinates in a rectangle from the plane, *Taiwanese Journal of Mathematics*, 4, 775-788, (2001).
- [11] Fejér, L., Über die Fourierreihen, II, *Math. Naturwiss. Anz Ungar. Akad., Wiss*, 24, 369-390(1906), (in Hungarian).
- [12] Gorenflo, R., Mainardi, F., *Fractional calculus: integral and differential equations of fractional order*, Springer Verlag, Wien, 223-276, (1997).
- [13] Hadamard, J., Etude sur les propriétés des fonctions entières et en particulier d'une fonction considérée par Riemann, *J. Math. Pures. et Appl.* 58, 171-215, (1893).
- [14] scan, I., Hermite-Hadamard-Fejer type inequalities for convex functions via fractional integrals, 2014, arXiv:1404.7722v1.
- [15] Latif, M.A., Alomari, M., Hadamard-type inequalities for product two convex functions on the co-ordinates, *Int. Math. Forum*, 4(47), 2327-2338, (2009).
- [16] Latif, M.A., Alomari, M., On the Hadamard-type inequalities for h -convex functions on the co-ordinates, *Int. J. of Math. Analysis*, 3(33), 1645-1656, (2009).
- [17] Latif, M.A., Hussain, S., New inequalities of Ostrowski type for co-ordinated convex functions via fractional integrals, *Journal of Fractional Calculus and Applications*, Vol. 2, No. 9, pp. 1-15, (2012).
- [18] Miller, S., Ross, B., *An introduction to the fractional calculus and fractional differential equations*, John Wiley & Sons, USA, p.2, (1993).
- [19] Minculete, N., Mitroi, F-C., Fejer-type inequalities, *Aust. J. Math. Anal. Appl.* 9, no. 1, Art. 12, 8pp, (2012).
- [20] Özdemir, M.E., Set, E., Sarikaya, M.Z., New some Hadamard's type inequalities for co-ordinated m -convex and (α, m) -convex functions, *Hacettepe Journal of Mathematics and Statistics*, 40(2), 219-229, (2011).
- [21] Sarikaya, M.Z., On new Hermite Hadamard Fejér type integral inequalities, *Stud. Univ. Babeş-Bolyai Math.* 57 (3), 377-386, (2012).
- [22] Sarikaya, M.Z., Set, E., Ozdemir M.E., Dragomir, S.S., New some Hadamard's type inequalities for co-ordinated convex functions, *Tamsui Oxford Journal of Information and Mathematical Sciences*, 28(2), 137-152, (2012).
- [23] Sarikaya M.Z., Ogunmez, H., On new inequalities via Riemann-Liouville fractional integration, *Abstract and Applied Analysis*, Volume 2012, Article ID 428983, 10 pages, doi:10.1155/2012/428983.
- [24] Sarikaya, M.Z., Set, E., Yaldiz, H., Basak, N., Hermite-Hadamard's inequalities for fractional integrals and related fractional inequalities, *Mathematical and Computer Modelling*, 57, 2403-2407, (2013).
- [25] Sarikaya, M.Z., Ostrowski type inequalities involving the right Caputo fractional derivatives belong to L_p , *Facta Universitatis, Series Mathematics and Informatics*, Vol. 27 No 2, 191-197, (2012).
- [26] Sarikaya, M.Z., Yaldiz, H., On weighted Montgomery identities for Riemann-Liouville fractional integrals, *Konuralp Journal of Mathematics*, Volume 1 No. 1 pp. 48-53, (2013).
- [27] Sarikaya, M.Z., Yaldiz, H., On the Hadamard's type inequalities for L -Lipschitzian mapping, *Konuralp Journal of Mathematics*, Volume 1 No. 2 pp. 33-40 (2013).
- [28] Sarikaya, M.Z., On the Hermite-Hadamard-type inequalities for co-ordinated convex function via fractional integrals, *Integral Transforms and Special Functions*, Vol. 25, No. 2, 134-147, (2014).

Hatice YALDIZ received her Ph.D. degree in Applied Mathematics in 2016 from Düzce University, Düzce, Turkey. She has been in Western Kentucky University as a visiting researcher in 2015. Her major research interests include inequalities, Fractional theory, Discrete analysis, Time scales.

Mehmet Zeki SARIKAYA is a Professor at the Department of Mathematics in Düzce University, Düzce, Turkey. He received his Ph.D. degree in Applied Mathematics in 2007 from Afyon Kocatepe University, Afyon, Turkey. He is editor on the Konuralp Journal. He has a lot of awards. He has many research papers about the Theory of inequalities, Potential Theory, Integral equations and Transforms, Time-scales..

Zoubir DAHMANI is Professor, Labo. Pure and Appl. Maths., Faculty of Exact Sciences and Informatics, UMAB, University of Mostaganem, Algeria. He received his Ph.D. of Mathematics, USTHB (Algeria) and La ROCHELLE University (France), 2009. He has many research papers about the Evolution Equations, Inequality Theory, Fractional Differential Equations, Fixed Point Theory, Numerical Analysis, Probability Theory, Generalized Metric Spaces.

